

Near-Surface Circulation and Fate of Upper Layer Fresh Water from Rivers Runoff and Rain in the Bay of Bengal near Sri Lanka

Luca Centurioni
Scripps Institution of Oceanography
9500 Gilman Drive
Mail Code 0213
La Jolla, California 92103
phone: (858) 534-6182 email: lcenurioni@ucsd.edu

Award Number: N00014-13-1-0477

LONG-TERM GOALS

Improve the knowledge of the near-surface circulation in the BoB and of the pathways through which the freshwater fluxes occur.

OBJECTIVES

We want to collect direct observations of the seasonally reversing currents of the southwest BoB to improve the knowledge of the near-surface circulation and for the validation of numerical ocean circulation models. The science questions we to address include the descriptive effort of identifying the routes through which the export of fresh water can potentially occur. More specifically:

- 1) is the Sri Lanka Dome, which shows up very clearly in the summer geostrophic surface current maps (Figure 1) effective in maintaining a nearly closed recirculation cell east off Sri Lanka or do particles (drifters) quickly disperse and get entrained in the northward flowing EICC and westward flowing SMC?
- 2) Does the southward wintertime EICC flows around Sri Lanka and exports fresh water into the Arabian Sea or does it significantly retrofect eastward along the east coast of Sri Lanka, as some authors suggest?

APPROACH

To address the two questions above, we are deploying 3 SVP drifters/month off the coast of Sri-Lanka.

A large number of drifters will be deployed 2015 when an additional fleet of 30 salinity drifters will be used in conjunction of other IOP activities to map the SSS distribution and variability over the experiment area and to infer the surface freshwater export pathways from direct measurements.

The salinity drifters will also be important to validate the Aquarius SSS data, which will also be used to infer the SSS export routes in conjunction with the drifter data.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2014		2. REPORT TYPE		3. DATES COVERED 00-00-2014 to 00-00-2014	
4. TITLE AND SUBTITLE Near-Surface Circulation and Fate of Upper Layer Fresh Water from Rivers Runoff and Rain in the Bay of Bengal near Sri Lanka				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California San Diego, Scripps Institution of Oceanography, 9500 Gilman Drive, La Jolla, CA, 92093				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

WORK COMPLETED AND RESULTS

The deployment of drifters begun in September 2012. The first shipment of 15 drifters was delivered to Mr. Udaya Jinadasa of NARA in Colombo in July 2013. Mr. Jinadasa was trained on the SVP drifter deployment methodology and the first deployment of 3 drifters (2 Argos 3 and one Iridium) occurred on 9/27/2013 and was performed by the Sri Lanka Navy (Figure 1). To date, 67 drifters were deployed in support of ASIRI.




Figure 1: tracks of the drifter deployed since the beginning of the project, as of 11/13/2014.

Drifters were also provided in support of two ASIRI cruises in fall 2013 and summer 2014.

RESULTS

A synthesis of drifter/altimetry/wind data was performed. The maps of near-surface currents show a well-developed EICC off Sri Lanka, and meandering currents and mesoscale features in the south-central BoB. The EICC moved around Sri Lanka, and turned westward as it passed south of the island. The width of the EICC off the eastern side Sri Lanka was about 100 km. The results of the synthesis were compared with observations collected during a December 2013 cruise and were found to be in very good agreement. Such results are being further interpreted and prepared for a publication.

IMPACT/APPLICATIONS

- Capacity building in Sri Lanka
- The drifter data are posted in real-time to the GTS of the World Weather Watch

RELATED PROJECTS

The Global Drifter Program

PUBLICATIONS

Lucas, A. J., E. L. Shroyer, H. W. Wijesekera, H. J. S. Fernando, E. D'Asaro, M. Ravichandran, S. U. P. Jinadasa, *et al.* "Mixing to Monsoons: Air-Sea Interactions in the Bay of Bengal." *Eos, Transactions American Geophysical Union* 95, no. 30 (2014): 269-70.